

CONSTRUCTION OF *PASTEURELLA HAEMOLYTICA* VACCINES

TECHNICAL AREA OF THE INVENTION

The invention relates to the area of bacterial genetic engineering. In particular, it relates to the bacteria *Pasteurella haemolytica*.

BACKGROUND OF THE INVENTION

The microorganism *P. haemolytica* biotype A, serotype 1, is the principal causative agent of pneumonic pasteurellosis in cattle. If techniques could be developed for introducing exogenous DNA into *P. haemolytica*, it would be possible to produce site-specific mutations in this bacterium. Such mutants could provide "rationally" attenuated strains for use as live vaccines.

Attenuated auxotrophic mutants were first described by Bacon and Burrows in the early 1950's. They reported that attenuated auxotrophs of *Salmonella typhi* defective in the aromatic amino acid biosynthetic pathway were avirulent in mice. Subsequently, it has been demonstrated in widely diverse bacteria that disrupting the aromatic amino acid biosynthetic pathway produces attenuated organisms. For example, attenuated strains of the invasive bacteria *Salmonella typhi*, *Salmonella typhimurium*, *Shigella flexneri*, and *Yersina enterocolitica*, were generated by introducing mutations in their respective *aroA* genes. Also attenuation was produced in the non-invasive bacteria *Bordetella pertussis* and *Pasteurella multocida* through *aroA* inactivation. Strains which carry *aroA* mutations are unable to synthesize chorismic acid from which p-aminobenzoic acid, dihydrobenzoate, and aromatic amino acids are produced. It is likely that the

09210747 "131500
1047 "131500

absence of one or more of these compounds *in vivo* is responsible for the poor growth of *aroA* mutants in the hosts.

Live attenuated bacterial strains generally provide superior protection as compared to killed bacterial vaccines (bacterins). In general, live vaccines elicit a stronger cell mediated response in the host than do bacterins. The superior immunity provided by attenuated live organisms may be explained by their ability to induce expression of stress-proteins and, possibly, of certain toxins within the host. The immune response generated by live organisms would be directed against these abundant proteins and thereby provide better protection.

There is a long-felt and continuing need in the art for veterinary vaccines to protect cattle from *P. haemolytica* infection. There also is a need for techniques for introducing DNA into *P. haemolytica*.

SUMMARY OF THE INVENTION

It is an object of the invention to provide methods for mutagenizing *P. haemolytica*.

It is another object of the invention to provide a *P. haemolytica* gene for production of an enzyme for use in preparing genetic material for introduction into *P. haemolytica*.

It is yet another object of the invention to provide an enzyme for use in preparing genetic material for introduction into *P. haemolytica*.

It is still another object of the invention to provide a plasmid for unstable introduction of genetic material into *P. haemolytica*.

It is an object of the invention to provide *P. haemolytica* mutant strains.

It is another object of the invention to provide live, attenuated vaccines against *P. haemolytica* infection.

It is another object of the invention to provide genetically engineered *P. haemolytica*.

These and other objects of the invention are provided by one or more of the embodiments described below. In one embodiment of the invention a method for site-directed mutagenesis of *P. haemolytica* is provided. The method comprises the steps of: isolating a DNA region from *P. haemolytica* in which

region a mutation is desired; introducing a mutation into said DNA region to form a mutated DNA region; methylating said mutated DNA region with a methylating enzyme, to form methylated DNA, which methylated DNA is refractory to endonuclease cleavage at GATGC and GCATC sequences; introducing said methylated DNA into *P. haemolytica* to form transformants; and screening said transformants for those which have said mutation in said region on chromosomal DNA of said *P. haemolytica* cell.

In an alternative embodiment of the invention site-directed mutagenesis of *P. haemolytica* is accomplished by the steps of: isolating a DNA region from *P. haemolytica* in which region a mutation is desired; introducing a mutation into said DNA region to form a mutated DNA region; introducing said methylated DNA into a *P. haemolytica* cell which does not express a *PhaI* restriction endonuclease, to form transformants; and screening said transformants for those which have said mutation in said region on chromosomal DNA of said *P. haemolytica* cell.

In another embodiment of the invention an isolated and purified gene is provided. The gene encodes *PhaI* methyltransferase.

In still another embodiment of the invention another isolated and purified gene is provided. The gene encodes *PhaI* restriction endonuclease.

In yet another embodiment of the invention a preparation of *PhaI* methyltransferase is provided. The preparation is free from *PhaI* restriction endonuclease.

In still another embodiment of the invention a preparation of *PhaI* restriction endonuclease is provided. The preparation is free from *PhaI* methyltransferase.

In another embodiment of the invention a chimeric plasmid is provided which is suitable for unstable introduction of genetic material into *P. haemolytica*. The plasmid comprises a 4.2 kb *P. haemolytica* plasmid encoding a streptomycin resistance determinant deposited at the American Type Culture Collection as Accession No. ATCC 69499; and a plasmid which cannot replicate in *P. haemolytica*.

In an additional embodiment of the invention a *P. haemolytica* mutant is provided. The mutant is made by the process of the invention described in more detail below.

In another embodiment of the invention a *P. haemolytica* mutant is provided which does not express the *PhaI* restriction endonuclease.

In another embodiment of the invention a *P. haemolytica aroA* mutant is provided.

In still another embodiment of the invention a vaccine is provided. The vaccine comprises an attenuated, live, mutant of *P. haemolytica* which has an *aroA* mutation.

In yet another embodiment of the invention an isolated and purified *P. haemolytica* strain is provided. The strain has been genetically modified by the introduction of DNA.

These and other embodiments of the invention provide the art with the means to construct desirable mutants of the economically important and previously intractable pathogen *P. haemolytica*.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Determination of *PhaI* cleavage positions alongside that of *SfaNI*. Lanes 1 and 3 cut with *PhaI*; lanes 2 and 4 cut with *SfaNI*. The cleavage products of *PhaI* and *SfaNI* migrated 0.5 bp faster than the corresponding sequence bands because the labeled primer for extension had a 5' phosphate, whereas the primer for sequencing did not (Juarin et al., *Gene* 39:191-201 (1985)).

Figure 2. Protection against *PhaI* digestion by cloned *PhaI*-methyltransferase. Lanes 1 and 2 plasmid pPh Δ AroACm^R-pD80 from *E. coli* DH10B incubated without and with *PhaI*. Lanes 3 and 4 plasmid pPh Δ AroACm^R-pD80 from *E. coli PhaIMtase* incubated without and with *PhaI*.

Figure 3. Southern blot analysis of *P. haemolytica* strain NADC-D60 DNA digested with *EcoRI* lane 1, *ClaI* lane 2, *PstI* lane 3, or *HindIII* lane 4. The membrane was hybridized with an *E. coli aroA* probe and washing was performed under low-stringency conditions.

6

(See also: 1) (See ID No. 2)
Figure 4. Nucleotide sequence_A and deduced amino acid sequence_A of *P. haemolytica aroA*.

Figure 5. Construction of a *P. haemolytica aroA* mutant. The hybrid plasmid pPharoA⁻Amp^RpD70 was successfully used to produce an *aroA* mutant.

Figure 6. Southern hybridization of genomic DNAs from the parental strain, *P. haemolytica* strain NADC-D60, the *aroA* mutant, and *P. haemolytica* strain NADC-D70 and the hybrid plasmid pPharoA⁻Amp^RpD70. All the DNAs used in the blots shown here were digested with *Hind*III. Figure 6A. Lanes: 1, *P. haemolytica* strain NADC-D60; 2, *aroA* mutant; 3, pPharoA⁻Amp^RpD70 probed with *P. haemolytica aroA*. Figure 6B. Lanes: 1, *P. haemolytica* strain NADC-D60; 2, *aroA* mutant; 3, pPharoA⁻Amp^RpD70 probed with *P. haemolytica* Amp^R fragment. Figure 6C. Lanes: 1, *P. haemolytica* strain NADC-D70; 2, *aroA* mutant; 3, pPharoA⁻Amp^RpD70 probed with *P. haemolytica* Amp^R plasmid. Figure 6D. 1, *P. haemolytica* strain NADC-D60; 2, *aroA* mutant; 3, pPharoA⁻Amp^RpD70 probed with pBCSK. DNA was isolated from *P. haemolytica* strain NADC-D70 and run in Lane 1 of blot B to demonstrate that if plasmid DNA was present in the bacteria it would also be present in our DNA preparations.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

It is a discovery of the present invention that *P. haemolytica* contains at least one restriction-modification system, called herein the *PhaI* system. Both the restriction endonuclease and the methyltransferase have been molecularly cloned. One such molecular clone (*E. coli PhaIMtase*) has been deposited at the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, 20852, USA, on December 2, 1993, under the terms of the Budapest Treaty as Accession No. ATCC 69500. A preliminary sequence of the methyltransferase gene has been determined. The predicted amino acid sequence of the methyltransferase contains sequence motifs which are consistent with an adenine-methylating specificity.

Provided with the molecular clone of *PhaIMtase* (Accession No. ATCC 69500) one of ordinary skill in the art can readily obtain a preparation of either or both enzymes free of other *P. haemolytica* proteins. A lysate of a non-*P. haemolytica* bacterium carrying one of the cloned enzymes would provide such a

10

20

30

preparation. If one desires a preparation of each of the enzymes free of the other enzyme, one of skill in the art can readily subclone to separate the two genes. The methyltransferase gene has been cloned into a plasmid which when introduced into a cell produces *PhaI* methyltransferase but is free of the *PhaI* restriction endonuclease. The *PhaI* restriction endonuclease gene can be cloned on a plasmid free of the methyltransferase gene by introduction of the cloned gene into host cells which express either the *PhaI* or the *SfaNI* methyltransferase.

Provided with *PhaIMtase* (ATCC Accession No. ATCC 69500) one of skill in the art can also readily obtain an isolated and purified gene encoding either or both the *PhaI* restriction and methyltransferase enzymes. Standard techniques, such as cesium chloride gradients, phenol and chloroform extractions, etc., can be used for purifying plasmid DNA from the deposited *E. coli* bacteria. The genes can be isolated together from the deposited bacteria, or they can be subcloned, as discussed above, to isolate the two genes from each other.

It has also been discovered by the present inventors, that a barrier to transformation of *P. haemolytica* can be overcome by treating DNA with a methylating enzyme, such as the *PhaI* methyltransferase. Such enzymes modify DNA substrates such that endonucleases which recognize 5'-GATGC-3' or 5'-GCATC-3' sequences are inhibited in their ability to digest such modified substrates. Examples of such endonucleases are *PhaI* endonuclease and *SfaNI* endonuclease. While applicants do not wish to be bound by any particular hypothesis on the mechanism of action of such methyltransferase enzymes, it appears that the *PhaI* methyltransferase methylates specific adenine residues in DNA.

Methylation of DNA substrates for transformation (electroporation, or other means of introduction of DNA into cells) can be accomplished *in vitro* or *in vivo*. For *in vitro* methylation, DNA is incubated with a preparation of methyltransferase in the presence of a methyl donor, such as S-adenosylmethionine (SAM). *In vivo* methylation can be accomplished by passaging the DNA substrate through a bacterium which contains an appropriate methyltransferase, such as *PhaI* or *SfaNI* methyltransferase. A mutant or natural variant of *P. haemolytica* which lacks the

PhaI endonuclease could also be used to prepare DNA for subsequent introduction into *P. haemolytica*. Such a mutant can be made *inter alia* according to the method for site-directed mutagenesis disclosed herein.

Site-directed mutagenesis of *P. haemolytica* can be accomplished according to the present invention by first isolating a wild-type DNA region from *P. haemolytica*. As described below in the examples, an *aroA* gene can be isolated using *aroA* DNA from other bacteria as hybridization probes. The sequence of the *P. haemolytica aroA* gene is shown in SEQ ID NO. 1. Similarly other genes can be isolated from *P. haemolytica*. Another desirable gene for mutations is the *PhaI* endonuclease gene, which is provided in *PhaIMtase* (ATCC Accession No. ATCC 69500). Other genes in which mutations may be desirable are genes in the leukotoxin operon (C, A, B, D) and neuraminidase. A mutation is created in the isolated, wild-type DNA region according to any method known in the art. For example, the isolated DNA can be chemically mutagenized, either in a bacterium or *in vitro*. Alternatively, restriction endonucleases can be used to create precise deletions or insertions *in vitro*. Other methods as are known in the art can be used as is desirable for a particular application.

After *P. haemolytica* DNA has been isolated and mutagenized, it is methylated as described above. Then it can be introduced into *P. haemolytica* according to any technique known in the art, including but not limited to transfection, transformation, electroporation, and conjugation. Alternatively, rather than methylating the mutagenized DNA and introducing it into a *P. haemolytica* which expresses *PhaI* restriction endonuclease, one can omit the methylation of the mutagenized DNA and introduce the mutagenized DNA into a *P. haemolytica* cell which does not express the *PhaI* restriction endonuclease. Such cells can be isolated from nature by extensive screening, isolated following chemical mutagenesis of a cell which does express the *PhaI* restriction endonuclease, or made by the site-directed mutagenesis method disclosed herein.

According to one aspect of the invention, the mutagenized and methylated *P. haemolytica* DNA region is introduced into a *P. haemolytica* cell on a plasmid which includes a *P. haemolytica* approximately 4.2 kb streptomycin resistance

determining plasmid (pD70). This plasmid has also been deposited at the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, 20852, USA, on December 2, 1993, under the terms of the Budapest Treaty as Accession No. ATCC 69499. While applicants do not wish to be bound by any particular theory, it appears that the pD70 streptomycin resistance determining plasmid allows the introduced DNA to be replicated and maintained, albeit unstably, for a period of time sufficient to allow gene conversion (replacement of the chromosomal copy of the gene with the introduced mutant copy of the gene) to occur. Gene conversion can be monitored *inter alia* by Southern hybridization with probes to the gene of interest, by screening for genetic markers on the introduced DNA construct (such as ampicillin^R or streptomycin^R), and by screening for the presence/absence of plasmid in the transformed cells' progeny.

A chimeric plasmid, as described above, is provided which is suitable for the unstable introduction of DNA into *P. haemolytica*. The chimeric plasmid comprises the approximately 4.2 kb streptomycin resistance determining plasmid, pD70, as well as a plasmid which cannot replicate in *P. haemolytica* but can replicate in another cell type. To use such a chimeric plasmid, typically a region of the chromosome of *P. haemolytica* which has been mutagenized is ligated into the plasmid. Maintenance of the chimeric plasmid in *P. haemolytica* can be selected, for example by using an appropriate antibiotic to which the plasmid confers resistance. After a selected number of generations, antibiotic selection can be removed, and the cells tested to determine whether the introduced region of *P. haemolytica* has replaced the genomic copy.

Also provided by the present invention are mutant strains made by the disclosed method of site-directed mutagenesis. One such mutant (NADC-D60 *aroA*⁻) has been deposited at the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, 20852, USA, on December 2, 1993, under the terms of the Budapest Treaty as Accession No. ATCC 55518. Such mutants can provide the veterinary arts with attenuated, live strains of *P. haemolytica* which are suitable for vaccines to induce protective immunity against *P. haemolytica* infection. For vaccine production, it is desirable that the mutation which

attenuates the *P. haemolytica* be an essentially non-reverting mutation. Typically these are deletion or insertion mutations, the latter not being caused by a transposable element. Strains which contain multiple attenuating mutations may also be used, so that the risk of reversion to a wild-type, virulent *P. haemolytica* is vanishingly small.

Another mutant strain which can be made by the site-directed mutagenesis method disclosed is one which is *PhaI* restriction endonuclease negative. Such a strain is useful for genetic engineering in *P. haemolytica*. Such a strain can be a recipient of DNA which is not *PhaI* methyltransferase methylated, yet would yield DNA which is *PhaI* methyltransferase methylated.

The present invention thus allows those of ordinary skill in the art to stably introduce DNA into *P. haemolytica*. The DNA can be from other strains or species. The DNA can be artificially modified or in its native state. If recombination into the genome is desired two regions of flanking homology are preferred. Such techniques are generally known for other bacteria, but have been hitherto unsuccessful in *P. haemolytica* due to its restriction system.

Vaccines are typically formulated using a sterile buffered salt solution. Sucrose and/or gelatin may be used as stabilizers, as is known in the art. It is desirable that the *P. haemolytica* vaccines of the invention be administered by the intranasal or intratracheal route, but subcutaneous, intramuscular, intravenous injections also may be used. Suitable formulations and techniques are taught by Kucera U.S. 4,335,106, Gilmour U.S. 4,346,074, and Berget U.S. 4,957,739. Typically, between 10^7 and 10^{11} CFU are administered per dose, although from 10^5 to 10^3 CFU can be used. Adjuvants also may be added.

EXAMPLES

Example 1

This example demonstrates the isolation and characterization of the type II_s restriction endonuclease *PhaI*.

Bacterium, growth, and crude extract

Pasteurella haemolytica serotype 1, strain NADC-D60, was grown 16 hours on 4 Columbia blood agar base plates (100ml total volume, Difco, Detroit, MI) without supplemental blood. The cells were harvested in TE (10mM Tris, 1mM EDTA, pH 8.0), pelleted by centrifugation at 16,000 G for 5 minutes at 4°C, and washed once in TE. The washed pellet was resuspended in 1.5 ml chromatography running buffer (20mM NaPO₄, 10mM 2-mercaptoethanol, pH 7.5, 4 C) and placed on ice. The bacterial cells were disrupted by sonication for 2 minutes in 15 second bursts. Debris and unbroken cells were removed by centrifugation at 16,000 G for 10 minutes and then filtration of supernatant through a 0.45 um HA membrane. No further treatment of the crude extract was performed prior to chromatography.

Chromatographic separation of proteins

All chromatographic procedures were performed at room temperature. Prepacked heparin-sepharose columns [Econopac heparin columns, Bio-Rad, Richmond, CA] were equilibrated as recommended by the manufacturer. A flow rate of 0.5 ml/minute was used for separation, controlled by 2 HPLC pumps and a controller [Beckman Instruments, Inc, Fullerton, CA]. One ml of crude extract was injected and 10 ml of running buffer was used to wash the column. A linear gradient from 0 to 0.5 M NaCl in 60 ml of running buffer was used to elute proteins. The column was washed with 2M NaCl in running buffer at 2.0 ml/minute as recommended by the manufacturer, then re-equilibrated to initial conditions of 0 M NaCl in running buffer prior to additional runs. Fractions (1.0 ml) were stored on ice prior to activity assay, then frozen at -20°C.

Assay for restriction endonuclease activity

Aliquots, 5 µl, of the chromatographic fractions were incubated with 1 µl 12 mM MgCl and 0.25 µg un methylated bacteriophage lambda DNA (New England Biolabs) at 37°C for 2 hours. After addition of tracking dye, and electrophoresis on a 1% agarose gel in TBE buffer, the banding patterns were visualized by ethidium bromide staining and UV illumination. The active fractions (6ml) were pooled, concentrated 10-fold on 30,000 MW cutoff ultrafilters, and

brought to final concentrations of 150 mM NaCl, 10 mM NaPO₄, 0.1 mM EDTA, 5 mM 2-mercaptoethanol, 0.25 µg/ml BSA, and 50:50 vol:vol glycerol [pH 7.5] for storage at -20°C.

Determination of the recognition sites for *Pha* I

The recognition sequence was identified using digestion of pBluescript (Stratagene, LaJolla, CA), which resulted in 4 fragments of approximate size 1476, 1057, 252, and 184 base pairs. Double digestion with *Pha*I and either *Xho*I or *Sac*I, which cut at opposite ends of the polylinker, showed that one *Pha*I site mapped at approximately nucleotide 1245, and another at 2735. Additional double digestions with *Ava*II, *Bgl*III, *Dra*II, *Pvu*I and *Sca*I were used to map the remaining 2 *Pha*I sites at approximately nucleotides 2300 and 2490, consistent with the sequences 5'-GATGC-3' and 5'GCATC-3'. Further confirmation was made with *Pha*I digests of ΦX174 and pUC19 DNA, and by sequencing pBluescript *Pha*I fragments filled in and cloned into pBluescript. Single-stranded ΦX174 DNA was digested to determine if *Pha*I has activity on this substrate.

Determination of the cleavage sites for *Pha* I

The cleavage site was identified by digestion of a primed-synthesis reaction on pBluescript derivatives (Brown et al. (1980) J. Mol. Biol. 140:143-148). An oligonucleotide containing the *Pha*I site was annealed and ligated with *Sma* I-cleaved pBluescript SK+ and SK-DNA. Single-stranded DNA containing each orientation was selected and used for the template. Four standard dideoxy DNA sequencing reactions were performed for each template with an appropriate primer. Additional reactions containing no dideoxy terminator were extended through the *Pha*I site with the Klenow fragment of DNA polymerase I using ³²P-endlabelled primer with both templates. The extension reaction was stopped by chloroform extraction followed by ethanol precipitation. *Pha*I or *Sfa* NI endonuclease was added to the additional reactions and allowed to digest the DNA for 2 minutes. The reaction was stopped by addition of gel loading buffer and heating to 80°C for 3 minutes.

A new restriction endonuclease, *Pha*I, an isochizomer of *Sfa*NI (Roberts (1990) Nucl. Acids Res. 18 (Suppl.), 2331-2365), was isolated from *Pasteurella*

haemolytica serotype 1, strain NADC-D60, obtained from pneumonic bovine lung. *PhaI* recognizes the 5 base non-palindromic sequence 5'-GCATC-3' and 5'-GATGC-3'. Cleavage occurs five bases 3' from the former recognition site and nine bases 5' from the latter recognition site.

Under our experimental conditions, endonuclease activity was eluted from heparin-sepharose columns by 275 to 325 mM NaCl. A single pass through these columns was sufficient to allow identification of both the DNA recognition specificity and cleavage site. Approximately 5000 units of *PhaI* per gram of wet cells were recovered. In contrast to *SfaNI*, optimal conditions for *PhaI* digestion required NaCl or KCl concentrations below 50 mM; > 50% reduction in activity was observed at the 100 mM NaCl optimum of *SfaNI*.

Digests of pBluescript resulted in 4 fragments of approximate size 1476, 1057, 252 and 184 bp. Double digestion with *PhaI* and either *XhoI* or *SacI* mapped 2 *PhaI* sites, one at approximately nucleotide 1245, and another at 2735 of pBluescript. Additional double digestions with *PhaI* and each of *AvaII*, *BglII*, *DraI*, *PvuI*, or *ScaI* mapped the remaining 2 *PhaI* sites at approximately nucleotides 2300 and 2490, consistent with the sequences 5'-GATGC-3' and 5'-GCATC-3'. Digests of pUC19, and Φ X174 confirmed the recognition specificity of 5'-GCATC-3', which is the same as that of *SfaNI*. Double digests of pBluescript with *PhaI* and *SfaNI* resulted in patterns identical to those using either enzyme alone. DNA containing the recognition sequence 5'-GATGC-3' cut 9 nucleotides 5' to the end of the recognition site with both *PhaI* and *SfaNI*. (Figure 1, lanes 1 and 2) DNA containing the recognition sequence 5'-GCATC-3' cut 5 nucleotides 3' to the end of the recognition site with both *PhaI* and *SfaNI*. (Figure 1, lanes 3 and 4)

β 5'...GCATCNNNNN↓NNNN...3' (SEQ ID NO:3)

β 3'...CGTAGNNNNN NNNN↑...5' (SEQ ID NO:4)

These data confirm that *PhaI* is a true isoschizomer of *SfaNI*. *PhaI* like *SfaNI* is a type IIs enzyme (Roberts, *Nucleic Acids Res.* 18:2331-2365 (1990)). The type IIs restriction enzymes, like the more common type II restriction enzymes, recognize specific sequences and cleave at predetermined sites. Type IIs enzymes,

however, neither recognize palindromic sequences nor cleave internally to the recognition sequence (Szybalski, *Gene* 100:13-26 (1991)).

Example 2

This example demonstrates the molecular cloning of *PhaI* endonuclease and methyltransferase.

Cosmid Library Construction

High-molecular weight DNA for cosmid cloning was prepared by the large scale DNA isolation method described for gram-negative bacteria in Ausabel et al. (*Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, NY, NY (1987)). Approximately 100 μ g of *P. haemolytica* strain NADC-D60 genomic DNA was digested with 100U of *ApoI* in NEB buffer #3 at 50°C for 10 minutes. Following digestion, the DNA was phenol-chloroform extracted and ethanol precipitated. The DNA was resuspended in 100 μ l TE and layered onto a linear gradient of 10-40% sucrose (Schwartz-Mann Ultrapure) in 10 mM Tris HCl, 1 mM EDTA, 100 mM NaCl, pH 8.0. After centrifugation in a SW40 (Beckman Inst.) at 20,000 RPM for 20 hr, gradient fractions were collected and restriction fragments of approximately 30 kb in length were ligated into *Eco* RI-digested calf alkaline phosphatase-treated cosmid vector pLAFRX (Ausabel, *supra*). A standard ligation mixture contained 1 μ g vector, 3 μ g *P. haemolytica* DNA and 5 Weiss U of T4 ligase in a volume of 10 μ l. The ligation mixture was incubated at 10°C for 16 hr. The DNA was packaged using Promega packaging extract (Promega, Madison, WI) according to the manufacturers' recommendations. *E. coli* HB101 transduced with the recombinant cosmid library were plated on 2XYT plates containing 10 μ g/ml tetracycline. Cloning efficiencies were approximately 10^4 recombinant colonies per μ g of genomic DNA.

Cloning of *PhaI* endonuclease and methyltransferase gene

Approximately 1 μ g of the recombinant *P. haemolytica* cosmid library was digested with *PhaI* restriction enzyme. The digested DNA was phenol-chloroform-isoamyl alcohol-extracted, ethanol precipitated, and resuspended in TE buffer. The DNA was electroporated into *E. coli* AP1-200-9 (Piekarowicz et al., *Nucl. Acids Res.* 19:1831-1835 (1991)) and the cells were plated on LB-broth plates

containing 20 $\mu\text{g/ml}$ tetracycline and 35 $\mu\text{g/ml}$ Xgal. The transformed cells were incubated at 42°C for 18 hours and transferred to 30°C for 4 hours. The cells were moved again to 42°C and blue colonies, indicating the presence of a cloned methyltransferase gene, were isolated and analyzed. The colonies were screened for restriction endonuclease activity by the technique of Schleif (*Method in Enzymology*, vol. 65, part I, pp. 19-23 (1980)). Double-stranded DNA mini-preps isolated from restriction endonuclease-positive colonies were analyzed for resistance to digestion by *PhaI*. Recombinant colonies resistant to *PhaI* digestion were presumed to contain a *PhaI* methyltransferase gene. Cosmid DNA from these cells was electroporated into *E. coli* DH10B (BRL, Gaithersburg, Maryland) and the cells were plated on LB-broth plates containing 20 $\mu\text{g/ml}$ tetracycline. The transformants containing the *PhaI* methyltransferase gene were designated *E. coli* strain *PhaIMtase*.

After digestion with *PhaI* and transformation of AP1-200-9 strain of *E. coli*, fifteen cosmid clones of *P. haemolytica* genomic DNA were tested for endonuclease activity. The nine clones which were endonuclease-positive were tested for *PhaI* methyltransferase activity. All nine expressed methyltransferase activity in addition to endonuclease activity, as evidenced by resistance to digestion by *PhaI* of genomic DNA recovered from transformed *E. coli*. The selective recovery of clones containing functional methyltransferase was due to previous digestion of the cosmid library with *PhaI* prior to transformation of *E. coli*. Recovery of clones containing both *PhaI* endonuclease and methyltransferase activity is not surprising since restriction and modification enzymes have previously been shown to be closely linked (the proximity of such genes has obvious implications to gene inheritance and the survival of the organism). The AP1-200-9 strain of *E. coli* (used to screen the cosmid library in this experiment) was designed by Piekarowicz et al., to give color selection for DNA-modifying enzymes (genes). The *mrr* and *mcr* systems, with a temperature-sensitive phenotype, induce inducible locus of the SOS response allows for color selection. All the transformants were blue after incubation at the permissive temperature for the *mcr/mrr* systems. Recovery of clones containing both *PhaI* endonuclease and

methyltransferase activity is not surprising since restriction and modification enzymes have previously been shown to be closely linked (the proximity of such genes has obvious implications to gene inheritance and to the survival of the organism). (Wilson et al., *Annu. Rev. Genet.* 25:585-627 (1991).)

Example 3

This example demonstrates the construction and methylation of a hybrid shuttle vector for introduction of DNA to *P. haemolytica*.

The following hybrid DNA construct was generated during attempts to introduce site-directed mutations into *P. haemolytica*. The *aroA* gene of *P. haemolytica*, contained on a *Hind*III-*Acc*I fragment of genomic DNA from strain NADC-D60, was ligated into the *Hind*III-*Acc*I site of pBluescript. A 700 bp fragment was excised from the coding region of the *aroA* gene by double digestion with *Nde*I and *Spy*I. Following digestion, the fragment ends were made blunt by treatment with the Klenow fragment of *E. coli* polymerase I and deoxynucleoside triphosphates. The deleted plasmid was excised from a 1% agarose gel and electroeluted. The eluted DNA, designated pPh Δ aroA2, was phenol-chloroform extracted and ethanol precipitated. The fragment was resuspended in TE buffer and ligated with the Cm^R gene isolated from pBR325. The Cm^R gene was excised from pBR325 by double digestion with *Aat* II and *Cla*I and the fragment was made blunt and purified by the above methods. The Cm^R fragment ligated with pPh Δ aroA2 was given the designation pPh Δ aroACm^R. Transformation of *E. coli* DH10B with pPh Δ aroACm^R conferred Cm^R to the bacterium.

The hybrid plasmid pPh Δ aroACm^RpD80 was constructed by ligating *Sma*I digested pPh Δ aroACm^R with *Sca*I digested pD80 (4.2 kb amp^R plasmid from *P. haemolytica* serotype 1 strain NADC-D80). The resultant hybrid plasmid, approximately 11 kb in size, contained a ColE1 and *P. haemolytica* ori, amp^R, and Cm^R.

For methylation, the hybrid plasmid was electroporated into *E. coli* strain DH10B with or without a cosmid containing cloned *Pha*I methyltransferase gene. Plasmid DNA was isolated and purified by CsCl gradient centrifugation. *Pha*I methyltransferase-treated hybrid plasmid was electroporated into *P. haemolytica*

strain NADC-D60 and then was reisolated by the above procedures. Plasmid DNA was reisolated from an ampicillin-resistant *P. haemolytica* transformant by the above procedures. The isolated plasmid DNA was tested for resistance to *PhaI* digestion as shown in Figure 2.

Example 4

This example demonstrates that methylated DNA, but not unmethylated DNA, is able to transform *P. haemolytica*.

Pasteurella haemolytica strain NADC-D60 was grown in 250 ml Columbia broth (Difco) 3 hours at 37°C with shaking to late logarithmic phase. The bacteria were centrifuged at 5000 G 15 minutes and the pellet resuspended in 272 mM sucrose at 0°C. The bacteria were washed 4 times in 272 mM sucrose with 5 minute centrifugation at 16,000 G and finally suspended at 50:50 vol:vol packed bacteria:272 mM sucrose on ice. Competent bacteria (100 µl) were mixed with 1 µg hybrid plasmid DNA (harvested from three sources: *E. coli* DH10B with methyltransferase (*PhaIMtase*); *E. coli* DH10B without methyltransferase; *P. haemolytica* NADC-D60) in 3 separate 1mm electroporation cuvettes (Bio-Rad), plus a fourth no-DNA control. The cells were quickly electroporated after addition of DNA (Bio-Rad Gene pulser) at 1500 V, 800 ohm, 25 uFd with resultant time constants ranging from 7.8 to 8.9 msec. Columbia broth (1ml, 0°C) was immediately added to the electroporated cells and the suspension was kept on ice approximately 10 minutes. The electroporated cells were allowed to recover at 37°C with gentle shaking for 1 hour, then broth containing 20 µg/ml ampicillin was added to bring the final ampicillin concentration to 10 µg/ml and the cells were incubated an additional hour at 37°C with shaking. Ten-fold dilutions were plated in duplicate onto blood agar plates containing 5% bovine blood and 10 µg/ml ampicillin. Undiluted cells electroporated with hybrid plasmid obtained from *E. coli* containing *PhaI* methyltransferase were plated on 2 µg/ml chloramphenicol after the first hour of recovery. Colonies were enumerated after overnight incubation at 37°C and representative colonies were checked for plasmid content.

Hybrid plasmid (pPh Δ aroACm^RpD80) passed through *E. coli* containing *PhaI* methyltransferase in a cosmid was able to transform *P. haemolytica* serotype 1. The hybrid plasmid was stably maintained through multiple passages under selective pressure. Whereas DNA not exposed to *PhaI* methyltransferase was unable to transform *P. haemolytica*, DNA methylated by *PhaI* methyltransferase in *E. coli* yielded 10³ transformants per μ g plasmid (Table 1). Plasmid DNA passed through *P. haemolytica* yielded 10⁵ transformants per μ g plasmid. This experiment demonstrates that the restriction-modification system of *PhaI* is important for introduction of exogenous DNA into *P. haemolytica* serotype 1.

The plating efficiency of transformants was 2 logs lower on chloramphenicol than on ampicillin. All transformants recovered, however, were resistant to both ampicillin, and chloramphenicol upon passage.

The possibility that a system similar to *E. coli mcr, mrr*, is active in *P. haemolytica* was investigated by passage of pPh Δ aroACm^RpD80 through *E. coli* strain GM2163 previously transformed with the recombinant cosmid containing *PhaI* methyltransferase (Raleigh et al., *Proc. Natl. Acad. Sci.* 83:9070-9074 (1986)). Since strain GM2163 is dam⁻, the resultant DNA would only be modified at *PhaI* sites (Marinus et al., *Mol. Gen. Genet.* 192:288-289 (1983)). Efficiency of transformation with this DNA, however, was not substantially different than that using DNA obtained from *PhaI* Mtase which is dam-methylated (Table 1). It is possible a second restriction system, not readily detectable in cell extracts, is active in *P. haemolytica* A1. Genes have been described in *Neisseria gonorrhoea* MS11 which encode for restriction enzymes which are expressed at levels too low to detect biochemically (Stein et al., *J. Bact.* 74:4899-4906 (1992)).

Table 1. Transformation efficiency of *P. haemolytica* NADC-D60 with hybrid plasmid pPhΔaroACm⁺pD80 purified from various sources^a.

Source of DNA ^b	Amp ^R transformants ^c CFU/μg DNA	Cm ^R transformants ^d CFU/μg DNA
<i>E. coli</i> DH10B	0	nd ^e
<i>E. coli</i> PhaIMtase	1x10 ³	5
<i>E. coli</i> GM2163	5x10 ²	nd
<i>P. haemolytica</i> NADC-D60	1x10 ⁵	nd

^a One μg DNA introduced by electroporation using same competent cell preparation.

^b Purified by CsCl-EtBr gradient centrifugation.

^c Colonies on plates containing 10 μg/ml ampicillin, cells recovered 2 hours prior to plating.

^d Colonies on plates containing 2 μg/ml chloramphenicol, cells recovered 1 hours prior to plating.

^e Not done.

This experiment demonstrates that the restriction-modification system of *PhaI* plays an important role in the difficulties researchers have encountered in their attempts to introduce exogenous DNA into *P. haemolytica* serotype 1. Protection against *PhaI* activity may allow genetic manipulation of this organism, which could lead to dramatic improvements in our understanding of pathogenesis and control of pneumonic pasteurellosis in cattle.

Example 5

This example demonstrates the molecular cloning and sequencing of *P. haemolytica* *aroA*.

Cloning of *P. haemolytica* *aroA*. Restriction fragments of *P. haemolytica* genomic DNA were fractionated by agarose gel electrophoresis. The fragments were probed for homology to a 1.3 kb *E. coli* *aroA* fragment by Southern analysis. Under conditions of low stringency, a 3.2-kb *HindIII* fragment of *P. haemolytica* genomic DNA hybridized with radiolabeled *E. coli* *aroA* (Fig. 3). The *HindIII* fragment was isolated from an agarose gel by electroelution and it was

cloned into *Hind*III digested pSK. The recombinant plasmid, pPharoA1, bearing *P. haemolytica aroA* was identified by complementation of *E. coli aroA* mutant AB2829 on M9 minimal media containing ampicillin. A *Cla* I, *Hind*III double digest of pPharoA1 generated a 2.2-kb fragment which was cloned into the *Acc*I and *Hind*III sites of pSK- giving rise to pPharoA2. Plasmid pPharoA2 also complemented growth of *E. coli* AB2829 on M9 minimal media. This plasmid was used to determine the sequence of *P. haemolytica aroA*.

Southern Blotting and Molecular Cloning of *P. haemolytica aroA*

Methods. *P. haemolytica* genomic DNA was prepared by the method for isolating DNA from gram-negative bacteria. Southern blotting of *P. haemolytica* restriction fragments fractionated by electrophoresis on a 0.75% agarose was performed as described previously. Blots were hybridized with a radioactively labeled 1.3-kb *E. coli aroA* fragment. The *aroA* probe was amplified (Gene-AMP by Perkin Elmer Inc., Branchburg, N.J.) from *E. coli* X-L1 Blue (Stratagene, Inc. S.D. CA) genomic DNA using PCR. The primers: 5' - TTCATGGAATCCCTTGACGTTACAACCCATC-3' and 5'-AGGCTGCCTGGCTAATCCGCGCCAG-3' used in the PCR reaction hybridize with *E. coli aroA* nucleotides -3 through 28 and 1308 through 1323 respectively. The primers were synthesized using an Applied Biosystems oligonucleotide synthesizer (Applied Biosystems Inc.) by the Nucleic Acids Facility, Iowa State University (Ames, IA). DNA was radiolabeled with [α - 32 P] dCTP using a random priming kit (Boehringer Mannheim Biochemicals, Indianapolis, IN). Nylon membranes (Hybond-N, Amersham Corp., Arlington Heights, IL) were incubated with hybridization solution 5X SSC (1X SSC is 0.15M NaCl and 0.015M sodium citrate), 5X Denhardt's solution (Maniatis, *Molecular Cloning*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1982)), 0.1% SDS, 10 μ g/ml sonicated salmon sperm DNA, containing 1×10^7 CPM of 32 P-labeled probe and 50% formamide at 42°C. After hybridization for 18 hours, membranes were washed twice with 1X SSC and 0.1% SDS for 10 minutes at RT and two times with 1X SSC and 0.1% SDS buffer at 42°C for 15 minutes. Membranes were exposed to X-AR (Eastman Kodak Co., Rochester, NY) at -80°C for 24 hours. A positive

signal corresponding to a 3.2-kb *Hind*III fragment of *P. haemolytica* chromosomal DNA was identified.

*Hind*III digested *P. haemolytica* DNA fragments ranging from 3.0-3.4 kb in length were electroeluted from a 1% agarose gel. The *Hind*III genomic fragments were added to *Hind*III digested alkaline phosphatase treated pSK-vector (Stratagene, Inc. S.D. CA) and ligated overnight at 10°C with T4 ligase (BRL, Gaithersburg, MD). The ligation mix was diluted 1:10 with distilled water and electroporated using a Gene Pulser (Bio-Rad Laboratories, Richmond, CA) into *E. coli aroA* mutant AB2829 (Pittard et al., *J. Bact.* 91:1494-1508 (1966)). A recombinant plasmid, pPharoA1, complemented AB2829 grown on M9 minimal media containing phosphate buffer, 1 mM MgSO₄, 0.1 mM CaCl₂, 0.2% glucose, thiamine 10 µg/ml, 1.5% Noble agar (Difco) and 50 µg/ml ampicillin (Ausubel et al., *supra*). A *Cla*I, *Hind*III double digest of pPharoA1 produced a 2.2 kb fragment which when cloned into the *Acc*I, *Hind*III sites of pSK- gave rise to pPharoA2. The recombinant plasmid, pPharoA2, which also complemented growth of AB2829 on minimal plates, was used to sequence *P. haemolytica aroA*.

Nucleotide sequence of *P. haemolytica aroA*. The nucleotide sequence and the deduced amino acid sequence of *P. haemolytica aroA* are shown in Fig. 4 and SEQ ID NOS: 1 and 2. An open reading frame of 1302 bases with a coding capacity of 434 amino acid residues was discerned. The deduced molecular weight is 47,296 and the G+C content of the *aroA* coding region is 43%. The predicted amino acid sequence of *P. haemolytica aroA* showed 75, 70, 69, and 68% identity with *Pasteurella multocida*, *Klebsiella pneumoniae*, *Yersenia enterocolitica*, and *Escherichia coli*, respectively.

P. haemolytica aroA, like *P. multocida aroA* (Homchampa et al. *Molec. Microbiol.* 23:3585-3593 (1992)), appears to be transcribed from its own promoter. This differs from the usual genetic arrangement in gram-negative bacteria where *aroA* and *serC* constitute an operon with *aroA* distal to the promoter. Evidence to support this claim are the findings that: (1) the nucleotide sequence upstream of *aroA* on clone pPharoA2 shows no homology with *serC* genes and (2) complementation of *E. coli* AB2829 by *P. haemolytica aroA*

contained on the 2.2 kb fragment is independent of the fragment's orientation on the cloning vector.

DNA sequencing and Analysis. DNA sequencing was done by the dideoxy nucleotide termination method with single or double stranded templates using the Sequanase 2.0 kit (United States Biochemicals, Cleveland, OH). A series of ordered deletions were made in *P. haemolytica aroA* contained on pPharoA2 using an Erase-a-base kit (Promega Corp. Madison, WI). Gaps in the sequence were completed using DNA primers synthesized by the DNA core facility at Iowa State University (Ames, IA). DNA sequence analysis was done with MacDNASIS Pro (Hitachi Software Ltd., San Bruno, CA) and MacVector (Kodak Co., New Haven, CT) software.

Example 6

This example demonstrates the construction of a defined *P. haemolytica aroA* mutant.

Construction of a *P. haemolytica aroA* mutant. The deletion plasmid, pPhΔaroACm^R (Table 2), was constructed from pPharoA2 as described above and amplified in *E. coli* containing a cosmid clone carrying the *PhaI* methyltransferase gene on a 20-kb *P. haemolytica* DNA fragment. Although resistant to *PhaI* endonuclease digestion, introduction of pPhΔaroACm^R into *P. haemolytica* strain NADC-D60 by electroporation failed to generate Cm resistant colonies. The inability to transform *P. haemolytica* with pPhΔaroACm^R suggested that plasmids containing a ColE1 origin do not replicate in this bacterium.

Table 2. Bacterial strains and plasmids used

Strains	Characteristics	Source/Reference
<i>E. coli</i>		
AB2829	K-12 strain with mutation in <i>aroA</i>	Pittard and Wallace (1966)
DH10B	Cloning strain used in this work	BRL
XL 1-Blue	Strain used for DNA sequencing	Stratagene
<i>P. haemolytica</i>		
NADC-D60	Serotype 1 plasmidless	NADC/R. Briggs
NADC-D70	Serotype 1 containing pD70	NADC/R. Briggs
NADC-D80	Serotype 1 containing pD80	NADC/R. Briggs
Plasmids		
pSK	cloning vector (Amp ^R)	Stratagene
pBCSK	cloning vector (Cm ^R)	Stratagene
pD70	4.2 kb plasmid encoding streptomycin ^R	NADC/R. Briggs
pD80	4.2 kb plasmid encoding Amp ^R	NADC/R. Briggs
pPharoA1	3.2 kb <i>Hind</i> III fragment containing <i>P. haemolytica aroA</i> (pSK)	This work
pPharoA2	<i>Hind</i> III <i>Cla</i> I digest of pPharoA1 resulted in 2.2 kb <i>aroA</i> fragment (pSK)	This work
pPharoA3	same insert as pPharoA2 on pBCSK	This work
pPhΔaroACm ^R	<i>Sty</i> I <i>Nde</i> I digest of pPharoA2 Cm ^R fragment inserted into deletion site	This work
pPhΔaroACm ^R pD80	<i>Sma</i> I digested pPhΔaroACm ^R joined to <i>Sca</i> I digested pD80	This work
pPhAmp ^R	2.2 kb <i>Sau</i> 3A fragment of pD80 cloned into pBCSK	This work
pPharoA ^R Amp ^R	Amp ^R fragment of pD80 inserted into unique <i>Nde</i> I site of pPharoA3	This work
pPharoA ^R Amp ^R pD70	<i>Hind</i> III digested pPharoA ^R Amp ^R joined to <i>Hind</i> III digested pD70	This work

Since we have shown that the *PhaI* methylated hybrid plasmid consisting of plasmids pPh Δ aroACm^R joined with *P. haemolytica* pD80 (Amp^R) could be used to transform *P. haemolytica* strain NADC-D60 (see above), we investigated the possibility that *aroA* mutants might arise after transformation with the hybrid plasmid by recombination with the genomic copy of the *aroA* gene, i.e., "replacement" of the gene. *P. haemolytica* harboring the hybrid plasmid pPh Δ aroACm^RpD80 were passed for > 100 generations in Columbia broth without antibiotics and plated onto blood-agar plates. The colonies were then replica plated onto blood-agar plates containing 5 μ g/ml ampicillin. All colonies had an Amp^R phenotype, suggesting that the plasmid was stable in *P. haemolytica*. This was confirmed by Southern blot analysis which showed that intact plasmid was present in all the Amp^R colonies that were analyzed.

Because the number of *P. haemolytica* transformants generated with hybrid plasmid pPh Δ aroACm^RpD80 (Amp^RCm^R) was 100-fold greater with plasmid isolated from *P. haemolytica* (10⁵ CFU/ μ g DNA) than from *E. coli* containing the *PhaI* methyltransferase gene (see above), we reasoned that a replacement plasmid isolated from *P. haemolytica* would be resistant to enzymatic digestion upon reintroduction into *P. haemolytica*, and thus more likely to give rise to mutants via homologous recombination. The improved efficiency is presumed to be the result of DNA modifications in *P. haemolytica* in addition to that of *PhaI* methylation.

With this in mind, hybrid plasmid pPh Δ aroACm^RpD80 was isolated from *P. haemolytica* strain NADC-D60 and CsCl purified by the methods described previously. The hybrid plasmid was digested with *Hind*III and *Xba*I to produce two fragments consisting of pD80 and pPh Δ aroACm^R. Linear deletion plasmid, pPh Δ aroACm^R, was isolated by electroelution and purified using "GlassMax" beads (BRL, Gaithersburg, MD). Approximately 5 μ g of the linear plasmid was electroporated into *P. haemolytica* NADC-D60. The cells were recovered in 1 ml Columbia broth and shaken at 37° C for 1 hour prior to plating on Blood-agar plates containing 10 μ g/ml chloramphenicol. No Cm^R colonies were detected after incubation at 37°C for 48 hours. However, this result was not totally unexpected

since there have been few reports of the successful establishment of linear DNA into bacteria.

Five μg of linearized $\text{pPh}\Delta\text{aroACm}^{\text{R}}$, isolated from *P. haemolytica*, was treated with Klenow and deoxynucleoside triphosphates to produce blunt ends. The DNA was then ligated with T4 ligase overnight to form a circular replacement plasmid. The plasmid was phenol chloroform extracted, ethanol precipitated, resuspended in distilled water, and reintroduced into *P. haemolytica* by electroporation. The cells were transferred to Columbia broth and allowed to recover for 1 hour. The cells were spread on blood-agar plates containing antibiotic and incubated at 37°C for 48 hours. This experiment also failed to generate Cm^{R} *P. haemolytica* colonies.

Additional efforts to produce an *aroA* mutant resulted in construction of a new replacement plasmid in which *aroA* was insertionally inactivated by the *P. haemolytica* β -lactamase gene. This antibiotic resistance cassette was chosen to select gene replacement candidates because we had found that survival of *P. haemolytica* transformed with $\text{pPh}\Delta\text{aroACm}^{\text{R}}\text{pD80}$ was approximately 100-fold greater (10^3 CFU/ μgDNA) on blood-agar plates containing ampicillin than on blood-agar plates containing chloramphenicol.

Molecular cloning of *P. haemolytica* β -lactamase gene was done as follows. Purified pD80 was partially digested with *Sau3A*, phenol-chloroform extracted, and ethanol precipitated. The fragments were resuspended in T.E. and ligated overnight into *Bam*HI-digested pBCSK (Stratagene Inc., La Jolla, CA). The ligated mixture was diluted 1:10 with water and electroporated into *E. coli* DH10B. The cells were recovered in 1 ml SOC for 1 hour and spread on LB-plates containing 50 $\mu\text{g/ml}$ ampicillin and 20 $\mu\text{g/ml}$ chloramphenicol. Restriction enzyme analysis on plasmid isolated from an ampicillin, chloramphenicol resistant *E. coli* clone revealed a 2.2 kb *P. haemolytica* insert in pBCSK. This plasmid was designated pPhAmp^{R} . To demonstrate that pPhAmp^{R} did not possess the pD80 origin of replication, the plasmid was amplified in *E. coli* DH10B which also contained the *PhaI* methyltransferase clone. Plasmid pPhAmp^{R} was isolated from *E. coli* as described previously, CsCl purified and introduced

into *P. haemolytica* by electroporation. Since this plasmid did not confer ampicillin resistance to *P. haemolytica* strain NADC-D60, we concluded that the antibiotic resistant fragment did not contain the pD80 origin of replication and that the fragment encoding the β -lactamase gene could be used to construct a deletion plasmid.

Construction of the deletion plasmid involved the following. The β -lactamase gene was excised from pPhAmp^R by *Hind*III, *Xba*I digestion and treated with Klenow and deoxy-ribonucleotides to generate blunt ends. The β -lactamase gene was ligated into the Klenow treated unique *Nde*I site of pPharoA3 (Fig. 5) to produce pPharoA⁻Amp^R. Insertional inactivation of *aroA* on pPharoA⁻amp^R was demonstrated by failure of the plasmid to complement AB2829. Plasmid pPharoA⁻Amp^R was amplified in *E. coli* DH10B (BRL) which also contained the recombinant cosmid carrying *PhaI* methylase recombinant cosmid. Although *PhaI* methylated pPharoA⁻Amp^R was resistant to digestion by *PhaI*, introduction of this plasmid into *P. haemolytica* failed to generate ampicillin resistant colonies.

To increase the likelihood of allelic replacement between the deletion plasmid's inactivated *aroA* and *P. haemolytica* chromosome, we constructed an *aroA*⁻ mutant-hybrid plasmid consisting of pPharoA⁻Amp^R and a 4.2-kb *P. haemolytica* plasmid (pD70, which confers streptomycin resistance (Sm^R)) (Fig. 5). The Sm^R plasmid was isolated from *P. haemolytica* using methods described previously. The str^R plasmid was digested at a unique *Hind*III site and ligated with *Hind*III digested pPharoA⁻Amp^R. The resultant hybrid plasmid, pPharoA⁻Amp^RpD70 (Fig. 5), was *PhaI* methyltransferase modified in *E. coli* DH10B containing the cosmid clone of the *PhaI* methylase gene. The hybrid plasmid was isolated from *E. coli*, CsCl purified and introduced into *P. haemolytica* strain NADC-D60 by electroporation. The cells were resuspended in Columbia broth for 2 hours at 37°C and spread on blood-agar plates containing 10 μ g/ml ampicillin. Transformation efficiency of the hybrid plasmid yielded approximately 10¹ ampicillin resistant colonies/ μ g DNA. Eight Amp^R colonies were grown overnight in Columbia broth containing 1 μ g/ml ampicillin. Chromosomal DNAs from the parental strain and from the Amp^R colonies were digested with *Hind*III and probed

by Southern blotting with *P. haemolytica* *aroA*, pBCSK, and pD70. The results indicated that intact pPharoA⁻Amp^RpD70 was present in the Amp^R colonies.

Eight Amp^R clones were grown overnight in Columbia broth containing 1 µg/ml ampicillin. Chromosomal DNAs from the parental strain and from the Amp^R clones were digested with *Hind*III and analyzed by Southern blotting with *P. haemolytica* *aroA*, pBCSK, and pD70 radio-labeled probes. The results indicated that intact pPharoA⁻Amp^RpD70 was present in the Amp^R clones (data not shown). The eight Amp^R cultures were transferred to Columbia broth containing 1 µg/ml ampicillin and cultured at 37° C. The bacteria were transferred to fresh media daily and this process was continued for approximately 100 generations. The eight cultures were streaked for isolation without antibiotic selection and a single colony of each was passed into Columbia broth containing either 1 µg/ml ampicillin or 1 µg/ml chloramphenicol. Two of the eight survived on the broth containing ampicillin, none on chloramphenicol. Passage from ampicillin broth onto blood-agar plates containing either ampicillin or chloramphenicol or streptomycin confirmed the two clones were Amp^R, Cm^S, Sm^S. Also the two Amp^R clones were spread onto plates of chemically-defined medium for *P. haemolytica* cultivation (Wessman, *Applied Microbiol.* 14:597-602 (1966)). This medium lacks the aromatic amino acid tryptophan. The parent strain grew on the defined medium but the Amp^R clones did not. Upon addition of tryptophan to the defined medium, growth of the Amp^R clones was comparable to that of the parent strain. The *E. coli* *aroA* mutant AB2829 also required tryptophan to grow on the chemically-defined medium for *P. haemolytica* cultivation. DNAs from the two colonies possessing Amp^R, Cm^S, Sm^S, *aroA*⁻ phenotypes were analyzed by Southern blotting. The results indicated that both had insertionally inactivated *aroA*s. Moreover, Southern blotting also confirmed that both pD70 and pBCSK sequences were no longer present in the *aroA* mutants (Figure 6).

Construction methods for *P. haemolytica* mutants. The 4.2 kb ampicillin resistance encoding plasmid of *P. haemolytica* (pD80) was partially digested with *Sau*3A and ligated into the *Bam*HI site of pBCSK⁺ (Cm^R) (Stratagene Inc., La Jolla, CA). The ligation mix was diluted 1:10 in distilled water and electroporated

into *E. coli* DH-10B (BRL, Gaithersburg, MD). After recovery in 1 ml SOC at 37°C, the cells were spread onto B-agar plates containing 50 µg/ml ampicillin. Plasmid, pPhAmp^R, contained a 2.2-kb *P. haemolytica* fragment which imparted ampicillin resistance to *E. coli* to up to 100 µg/ml. Plasmid, pPhAmp^R, was digested with *Hind*III and *Xba*I digestion and the fragment ends were made blunt by incubation with deoxynucleotide triphosphates and the large Klenow fragment of *E. coli* polymerase I. The fragment encoding ampicillin resistance was electroeluted. *P. haemolytica* *aroA* contained on pPharoA3 was digested at an unique restriction site within the coding region of *aroA* with *Nde*I and the fragment ends were made blunt as described previously. The fragment encoding ampicillin resistance was blunt-end ligated with T4 ligase into pPharo2 thus generating pPharoA-Amp^R. Plasmid pPharoA-Amp^R was digested with *Hind*III and dephosphorylated with calf alkaline phosphatase. A 4.2 kb plasmid encoding Sm^R isolated from *P. haemolytica* strain NADC-D70 (Chang et al., *J. DNA Sequencing and Mapping* 3:89-97 (1992)) was also digested with *Hind*III and the two plasmids were ligated with T4 ligase to generate the hybrid plasmid pPharoA-Amp^RpD70. The hybrid plasmid was electroporated into *E. coli* *Pha* IMtase which contained the *PhaI* methyltransferase gene on cosmid pLAFRX (Ausubel, *supra*).

P. haemolytica strain NADC-D60 is a plasmidless strain which was isolated from a cow with pneumonic pasteurellosis. The *PhaI* methylated hybrid plasmid was CsCl purified and 1 µg plasmid and 30 µl of *P. haemolytica* strain NADC-D60 were transferred to an 0.2 cm. cuvette and electroporated at 15,000 volts/cm with 800 ohms. The resultant time constant was approximately 9 milliseconds. Cells were transferred to 2 ml Bacto Columbia broth (Difco Labs, Detroit, MI) and incubated at 37°C for two hours and spread on Difco Columbia blood-agar plates containing 10 µg/ml ampicillin. Eight ampicillin resistant *P. haemolytica* colonies were isolated after incubation at 37°C for 18 hours. The colonies were then transferred to Bacto-Columbia broth containing 1 µg/ml ampicillin and incubated at 37°C. Daily passage into fresh medium containing 1 µg/ml ampicillin was carried out for three days at which time the cultures were transferred onto Columbia broth blood-agar plates containing 10 µg/ml ampicillin and incubated at

37°C overnight. The next day, colonies were replica-plated onto Columbia broth blood-agar plates containing 10 µg/ml or 50 µg/ml streptomycin and a chemically-defined medium for *P. haemolytica* cultivation (Wessman, *supra*). The defined medium contains 15 amino acids and includes the aromatic amino acids phenylalanine and tyrosine but not tryptophan. The clones unable to grow on the chemically-defined medium for *P. haemolytica* cultivation were presumed to be *aroA*⁻. Genomic DNA isolated from colonies with Amp^R, Cm^S, Sm^S, *aroA*-phenotypes were analyzed by Southern blotting. Southern blotting was performed as described previously with the exception that after hybridization the membranes were washed twice for 10 minutes each in 1x SSC and 0.1% SDS at 42° C and twice more for 15 minutes each in 0.1x SSC and 0.1% SDS at 65° C.

add
B'